

REFERENCES

- 1 Buehler JW, Berkman RL, Hartley DM, *et al.* Syndromic surveillance and bioterrorism-related epidemics. *Emerg Infect Dis* 2003;9:1197–204.
- 2 Dürheim DN, Speare R. Communicable disease surveillance and management in a globalised world. *Lancet* 2004;363:1339–40.
- 3 Pavlin JA, Mostashari F, Kortepeter MG, *et al.* Innovative surveillance methods for rapid detection of disease outbreaks and bioterrorism: results of an interagency workshop on health indicator surveillance. *Am J Public Health* 2003;93:1230–5.
- 4 Centers for Disease Control and Prevention. Syndromic surveillance: reports from a national conference, 2004. *MMWR* 2005;54(suppl).
- 5 Reingold A. If syndromic surveillance is the answer, what is the question? *Biosecure Bioterror* 2003;1:77–81.
- 6 Bravata DM, McDonald KM, Smith WM, *et al.* Systematic review: surveillance systems for early detection of bioterrorism-related diseases. *Ann Intern Med* 2004;140:910–22.
- 7 Centers for Disease Control and Prevention. Framework for evaluating public health surveillance systems for early detection of outbreaks: recommendations from the CDC working group. *MMWR Recomm Rep* 2004;53:1–11.
- 8 Heffernan R, Mostahari F, Das D, *et al.* Syndromic surveillance in public health practice, New York City. *Emerg Infect Dis* 2004;10:858–64.
- 9 Kulldorff M, Heffernan R, Hartman J, *et al.* A space-time permutation scan statistic for disease outbreak detection. *PLoS Med* 2005;2:e59.
- 10 Stoto M, Schonlau M, Mariano L. Syndromic surveillance: is it worth the effort? *Chance* 2004;17:19–24.
- 11 Kaufmann A, Pesik N, Meltzer M. Syndromic surveillance in bioterrorist attack. *Emerg Infect Dis* 2005;11:1487–8.

Speaker's corner

Autism and diet

Is exposure to antibodies and fatty acids in mother's milk associated with development of autism? Autism is a disease of unknown aetiology, but complex pathogenesis that seems to have become more common in recent decades. We suggest that efforts exploring the aetiology of autism should pay special attention to patterns of breast and artificial feeding in infancy.

Newschaffer grouped the aetiology of risks for autistic disorders into four groups: genetic predisposition of the mother, environmental factors acting on the mother, genetic predisposition of the child, and environmental factors affecting the child.¹ Historical incidence has been difficult to determine as the heterogeneous nature of the syndrome and the recent increase in awareness confound past estimates.² However, it is estimated that autism has increased threefold since the 1970s and that current rates are around 5–10 cases per 10 000 in the general population.¹ If prevalence is increasing it implies that even if there is a genetic component we also need to look for some environmental change that could be triggering any increase in prevalence.

There are well identified links between gastrointestinal disease and symptomatic autism. Wakefield claimed that diarrhoea occurred disproportionately in a control study of autistic children.³ Children with autism have unusual feeding patterns and in one study low appetite, narrow range of preferred dishes, and digestive autonomic nervous system reactions were significantly more common in the autistic group when compared with other children in one study.⁴

The postnatal growth of the human brain requires a range of fatty acids and the content of human milk changes with brain growth. By age 3 the brain of an autistic child undergoes a growth spurt and is larger than that of a normal

child at age 3.⁵ Tanoue and colleagues suggested that breast feeding may protect against autism. Researchers studied the weaning times of 145 autistic infants compared with a control group of 224 normal children. They found that the children in their control group breast fed significantly longer than the autistic infants. The researchers, who had previously found a correlation between incidence of autism and cyclical outbreaks of pneumonia and bronchiolitis, speculated that breast feeding lowered the risk of developing autism by protecting against pneumonia.⁶ Those with autism may be more vulnerable to infection during the newborn period. The evidence of links between diet and autism are suggestive and warrant further investigation.

M Potts, B Bellows

School of Public Health, University of California, Berkeley, USA
Correspondence to: Mr B Bellows, 314 Warren Hall, Berkeley, CA 94720 USA; bbellows@berkeley.edu

Conflicts of interest: none declared.

REFERENCES

- 1 Newschaffer C, Fallin D, Lee N. Heritable and nonheritable risk factors for autism spectrum disorders. *Epidemiol Rev* 2002;24:137–53.
- 2 Stokstad E. Epidemiology. Vaccine-autism link dealt blow. *Science* 2003;301:1454–5.
- 3 Wakefield AJ. Enterocolitis, autism and measles virus. *Mol Psychiatry* 2002;7(suppl 2):S44–6.
- 4 Lesinskiene S, Puras D, Kajokiene A, *et al.* [Features of nursing of the autistic children]. *Medicina (Kaunas)* 2002;38:412–19.
- 5 Courchesne E, Carper R, Akshoomoff N. Evidence of brain overgrowth in the first year of life in autism. *JAMA* 2003;290:337–44.
- 6 Tanoue Y, Oda S. Weaning time of children with infantile autism. *J Autism Dev Disord* 1989;19:425–34.